A service of the <u>U.S. National Library of Medicine</u> and the <u>National Institutes of Health</u>

My NCBI [Sign In] [Regi	2
[Sign In] [Regi	ster]

All Databases PubMed	Nucleotide Pr	otein Genome	Structure	OMIM PMC	Journals	Books	
Search PubMed	for		es à	Go Clear	Save Search		
Limits Preview/Index Histor	y Clipboard De	etails					
Display AbstractPlus	Show 20	Sort By	Send to				
All: 1 Review: 0		entering of the second	The second secon				
1: <u>Cell.</u> 1995 Feb 24;80(4):66	51-70.					Links	
The anticoagulation factor protein S and its relative, Gas6, are ligands for the Tyro 3/AxI family of receptor tyrosine kinases.				Related Links			
Stitt TN, Conn G, Gore M, Lai C, Bruno J, Radziejewski C, Mattsson K, Fisher J, Gies DR, Jones PF, et al.				Identification of Gas6, a putative ligand for Sky and Axl receptor tyrosine kinases, as a novel neurotrophic factor for hippocampal neurons[J Neurosci Res. 2002]			
Regeneron Pharmaceuticals, Incorporated, Tarrytown, New York 10591.				Indentification and regulation of receptor tyrosine kinases Rea and Mer and their ligand Gas6 in testicular som atic cells. [J Androl. 2000]			
We report the identification of ligands for Tyro 3 (alternatively called Sky, rse, brt, or tif) and Axl (alternatively, Ark or UFO), members of a previously orphan family of receptor-like tyrosine kinases. These ligands correspond to protein S, a protease regulator that is a potent anticoagulant, and Gas6, a protein related to			orphan cein S, a	Identification of Gas6 as a ligand for Mer, a neural cell adhesion molecule related receptor tyrosine kinase implicated in cellular transformation [Oncogene, 1997]			
protein S but lacking any known function. Our results are reminiscent of recent findings that the procoagulant thrombin, a protease that drives clot formation by cleaving fibrinogen to form fibrin, also binds and activates intracellular signaling				Identification of the product of growth arrest-specific gene 6 as a common ligand for Ad, Sky, and Mer receptor tyrosine kinases. [J Biol Chem. 1996]			
via a G protein-coupled cell surface receptor. Proteases and protease regulators that also activate specific cell surface receptors may serve to integrate coagulation with associated cellular responses required for tissue repair and growth, as well as to coordinate protease cascades and associated cellular			and	► Expression of the receptor protein-tyrosine kinases Tyro-3, AxI, and mer in the developing rat central nervous system. [J Comp Neurol. 2000]			
responses in other systems, such as those involved in growth and remodeling of the nervous system.				» See all Related Articles			
PMID: 7867073 [PubMed - indexed for ME	EDLINE]						
Display AbstractPlus	Show 20	Sort By	Send to				

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer